

GAUSS-MANIN CONNECTIONS AND LIE-RINEHART COHOMOLOGY

HELGE MAAKESTAD

ABSTRACT. In this note we prove existence of a generalized Gauss-Manin connection for a class of Ext-sheaves extending the construction in Katz paper [12]. We apply this construction to give a Gysin map for the K-groups of categories of coherent flat connections. We also give explicit examples of a generalized Gauss-Manin connection for a family of curves.

CONTENTS

Introduction	1
1. Gauss-Manin connections and Lie-Rinehart cohomology	2
2. Application: A Gysin map for K-theory of flat connections	7
3. Explicit examples: A Gauss-Manin connection for a family of curves	9
References	13

INTRODUCTION

In previous papers (see [17] and [18]) a theory of characteristic classes with values in Lie-algebra cohomology was developed: For an arbitrary Lie-Rinehart algebra L and any A -module W with an L -connection there exists Chern-classes (see [17])

$$c_i(W) \in \mathbf{H}^{2i}(L|_U, \mathcal{O}_U)$$

where U is the open subset where W is locally free, generalizing the classical Chern-classes with values in de Rham-cohomology. Moreover there exists a Chern-character (see [18])

$$ch : K(L) \rightarrow \mathbf{H}^*(L, A)$$

generalizing the classical Chern-character with values in de Rham-cohomology. The cohomology theory $\mathbf{H}^*(L, A)$ is a cohomology theory simultaneously generalizing de Rham-cohomology of A and Lie-algebra cohomology of L .

In this paper we consider the problem of constructing a Gauss-Manin connection for the cohomology group $\mathbf{H}^i(L, W)$ generalizing the Gauss-Manin connection existing for de Rham-cohomology. Consider the category of flat $\mathcal{L}$ -connections where

Date: Autumn 2005.

Key words and phrases. Lie-Rinehart algebras, Lie-algebra cohomology, Kodaira-Spencer maps, Gysin-maps, generalized Gauss-Manin connections, quotient singularities.

$\mathcal{L}$ is a quasi-coherent sheaf of Lie-Rinehart algebras on an arbitrary scheme X . Let $f : X \rightarrow S \rightarrow T$ be arbitrary maps of schemes, and let

$$\begin{array}{ccc} \mathcal{L} & \longrightarrow & \Theta_X \\ \downarrow F & & \downarrow df \\ f^*\mathcal{H} & \longrightarrow & f^*\Theta_S \end{array}$$

be any surjective map of Lie-Rinehart algebras. Let $\mathcal{K} = \ker(F)$. We construct a class of ext-sheaves giving rise to functors

$$\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(-, -) : \underline{\mathcal{L}\text{-conn}} \times \underline{\mathcal{L}\text{-conn}} \rightarrow \underline{\mathcal{H}\text{-conn}}.$$

hence we get for any pair of flat $\mathcal{L}$ -connections $\mathcal{V}, \mathcal{W}$ a canonical flat connection - the generalized Gauss-Manin connection:

$$\nabla : \mathcal{H} \rightarrow \underline{\mathrm{End}}_S(\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^*(\mathcal{V}, \mathcal{W})).$$

As an application we get in Theorem 2.2 Gysin-maps

$$F_! : K(\mathcal{L}) \rightarrow K(\mathcal{H})$$

defined by

$$F_![\mathcal{W}, \nabla] = \sum_{i \geq 0} (-1)^i [\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, \mathcal{W})]$$

between K-groups of categories of coherent flat connections.

We also construct in Theorem 3.2 a class of explicit examples of generalized Gauss-Manin connections on a Lie-Rinehart algebra $H \subseteq T_C$

$$\nabla_i : H \rightarrow \mathrm{End}_R(\mathbf{H}^i(L, W)).$$

associated to a family of curves $X \rightarrow C$, where X is a quotient surface singularity.

The aim of the construction is to use the Gysin-map to later on prove Riemann-Roch formulas for Chern-classes with values in Lie-Rinehart cohomology (see [15]).

1. GAUSS-MANIN CONNECTIONS AND LIE-RINEHART COHOMOLOGY

In this section we define Lie-Rinehart algebras, connections and generalized Lie-algebra cohomology and state some general properties of the constructions introduced. We also prove existence of a class of Ext-sheaves equipped with canonical flat connections - generalized Gauss-Manin connections.

First some definitions:

Definition 1.1. Let A be an S -algebra where S is a commutative ring with unit. A *Lie-Rinehart algebra* on A is a S -Lie-algebra and an A -module L with a map $\alpha : L \rightarrow \mathrm{Der}(A)$ satisfying the following properties:

$$\begin{aligned} (1.1.1) \quad & \alpha(a\delta) = a\alpha(\delta) \\ (1.1.2) \quad & \alpha([\delta, \eta]) = [\alpha(\delta), \alpha(\eta)] \\ (1.1.3) \quad & [\delta, a\eta] = a[\delta, \eta] + \alpha(\delta)(a)\eta \end{aligned}$$

for all $a \in A$ and $\delta, \eta \in L$. Let W be an A -module. An L -connection ∇ on W , is an A -linear map $\nabla : L \rightarrow \mathrm{End}_S(W)$ wich satisfies the *Leibniz-property*, i.e.

$$\nabla(\delta)(aw) = a\nabla(\delta)(w) + \alpha(\delta)(a)w$$

for all $a \in A$ and $w \in W$. We say that (W, ∇) is an L -module if ∇ is a homomorphism of Lie-algebras. The *curvature* of the L -connection, R_∇ is defined as follows:

$$R_\nabla(\delta \wedge \eta) = [\nabla_\delta, \nabla_\eta] - \nabla_{[\delta, \eta]}.$$

A Lie-Rinehart algebra is also referred to as a *Lie-Cartan pair*, *Lie-algebroid*, *Lie-pseudo algebra* or an *LR-algebra*. The notion of a Lie-Rinehart algebra and connection globalize to sheaves, hence we get a definition for any scheme X/S : It is a pair $(\mathcal{L}, \alpha)$ where

$$\alpha : \mathcal{L} \rightarrow \Theta_X$$

is a map of quasi-coherent sheaves satisfying the properties from Definition 1.1 locally. An $\mathcal{L}$ -connection is a map of quasi-coherent sheaves

$$\nabla : \mathcal{L} \rightarrow \text{End}_{\mathcal{O}_S}(\mathcal{W})$$

satisfying the criteria of Definition 1.1 locally.

Assume for the rest of the paper that only morphisms of schemes with well-defined map on tangent-spaces are considered.

We next define morphisms of LR-algebras (see [1] and [5]): Let $f : X \rightarrow S$ be a map of schemes. Let furthermore $\mathcal{L} \rightarrow \Theta_X$ and $\mathcal{H} \rightarrow \Theta_S$ be LR-algebras.

Definition 1.2. A *morphism* of LR-algebras, is a pair of maps (F, f) giving a commutative diagram

$$\begin{array}{ccc} \mathcal{L} & \xrightarrow{\alpha} & \Theta_X \\ \downarrow F & & \downarrow df \\ f^*\mathcal{H} & \xrightarrow{f^*\beta} & f^*\Theta_S \end{array}$$

Here F is a map of sheaves satisfying

$$(1.2.1) \quad F([x, y]) = [F(x), F(y)] + x.F(y) - y.F(x)$$

where $F(x) = \sum a_i \otimes x_i$ and $F(y) = \sum b_j \otimes y_j$ are local sections and the following holds:

$$x.F(y) = \sum x(b_j) \otimes y_j,$$

$$[F(x), F(y)] = \sum_{i,j} a_i b_j \otimes [x_i, y_j].$$

Let $\mathcal{U}(\mathcal{L}/\mathcal{H}) = f^*\mathcal{U}(\mathcal{H})$ be the *transfer-module* of the morphism F . Then $\mathcal{U}(\mathcal{L}/\mathcal{H})$ has a canonical $\mathcal{L}$ -connection.

Lemma 1.3. *The canonical $\mathcal{L}$ -connection on $\mathcal{U}(\mathcal{L}/\mathcal{H})$ is flat.*

Proof. Straight forward. □

In fact given any map $F : \mathcal{L} \rightarrow f^*\mathcal{H}$ it is easy to see that it is a map of LR-algebras if and only if the canonical connection on $\mathcal{U}(\mathcal{L}/\mathcal{H})$ is flat: flatness is equivalent to the equation 1.2.1. We get thus a category of LR-algebras, denoted LR-*alg.*

We define cohomology of LR-algebras as follows:

Definition 1.4. Let L be a Lie-Rinehart algebra and (W, ∇) a flat L -connection. The *standard complex* $C^\bullet(L, W)$ is defined as follows:

$$C^p(L, W) = \text{Hom}_A(L^{\wedge p}, W),$$

with differentials $d^p : C^p(L, W) \rightarrow C^{p+1}(L, W)$ defined by

$$d^p \phi(g_1 \wedge \cdots \wedge g_{p+1}) = \sum_{i=1}^{p+1} (-1)^{i+1} g_i \phi(g_1 \wedge \cdots \wedge \overline{g_i} \wedge \cdots \wedge g_{p+1}) + \sum_{i < j} (-1)^{i+j} \phi([g_i, g_j] \wedge \cdots \wedge \overline{g_i} \wedge \cdots \wedge \overline{g_j} \wedge \cdots \wedge g_{p+1}).$$

Here $g\phi(g_1 \wedge \cdots \wedge g_p) = \nabla(g)\phi(g_1 \wedge \cdots \wedge g_p)$ and overlined elements should be excluded. It follows that $C^\bullet(L, W)$ is a complex if and only if ∇ is flat. The cohomology $\mathbf{H}^i(C^\bullet(L, W))$ is denoted $\mathbf{H}^i(L, W)$ - the *Lie-Rinehart cohomology* of (W, ∇) .

In the litterature the standard complex is also referred to as the *Chevalley-Hochschild complex*. There exists a generalized universal enveloping algebra $U(L)$ with the property that there is a one to one correspondence between flat L -connections and left $U(L)$ -modules.

Proposition 1.5. *Let L be locally free. There is an isomorphism*

$$\mathbf{H}^i(L, W) \cong \text{Ext}_{U(L)}^i(A, W)$$

for all $i \geq 0$.

Proof. See [8]. □

Let for the rest of this section $f : X \rightarrow S$ be an arbitrary map of schemes defined over a base scheme T . Let $\mathcal{L} \rightarrow \Theta_X$ and $\mathcal{H} \rightarrow \Theta_S$ be LR-algebras, and let $F : \mathcal{L} \rightarrow f^*\mathcal{H}$ be a surjective map of LR-algebras with associated exact sequence

$$(1.5.1) \quad 0 \rightarrow \mathcal{K} \rightarrow \mathcal{L} \rightarrow f^*\mathcal{H} \rightarrow 0.$$

Proposition 1.6. *There is a functor*

$$F_! : \underline{\mathcal{L} - \text{conn}} \rightarrow \underline{\mathcal{H} - \text{conn}}$$

defined by

$$F_!(\mathcal{W}, \nabla) = (f_*(\mathcal{W}^\nabla), \overline{\nabla})$$

where $\mathcal{W}^\nabla$ is the sheaf of horizontal sections with respect to the induced connection on $\mathcal{K}$ and $\overline{\nabla}$ is a flat $\mathcal{H}$ -connection.

Proof. We assume $X = \text{Spec}(B)$, $S = \text{Spec}(A)$ and $T = \text{Spec}(R)$. The exact sequence in 1.5.1 becomes

$$0 \rightarrow K \rightarrow L \xrightarrow{F} B \otimes_A H \rightarrow 0.$$

Let W be a module with a flat L -connection ∇ and consider $W^\nabla \subseteq W$ the sub- A -module of sections with $\nabla(y)(w) = 0$ for all $y \in K$. We seek to define a canonical flat H -connection

$$\overline{\nabla} : H \rightarrow \text{End}_R(W^\nabla).$$

Let $x \in H$ and pick an element $y \in L$ with $F(y) = 1 \otimes x$. Define $\overline{\nabla}(x)(w) = \nabla(y)(w)$. This is well defined since $w \in W^\nabla$. We want to show we get an induced map

$$\overline{\nabla}(x) : W^\nabla \rightarrow W^\nabla.$$

Pick $h \in K$. We get

$$\begin{aligned}\nabla(h)\nabla(y)(w) &= \nabla(y)\nabla(h)(w) + \nabla([h, y])(w) = \\ &\nabla([h, y])(w).\end{aligned}$$

Since

$$F[h, y] = [Fh, Fy] + h.F(y) - y.F(h) = h.(1 \otimes x) = h(1) \otimes x = 0$$

it follows $[h, y] \in K$ hence $\nabla([h, y])(w) = 0$ and we have shown we get an induced map

$$\overline{\nabla}(x) : W^\nabla \rightarrow W^\nabla$$

hence we get an induced flat connection

$$\overline{\nabla} : H \rightarrow \text{End}_R(W^\nabla).$$

This argument globalize and the claim is proved. $\square$

We get thus in the situation where X and S are arbitrary schemes and $F : \mathcal{L} \rightarrow f^*\mathcal{H}$ surjective map of LR-algebras a left exact functor

$$F_! : \underline{\mathcal{L}\text{-conn}} \rightarrow \underline{\mathcal{H}\text{-conn}},$$

defined by

$$F_!(\mathcal{W}, \nabla) = (\mathcal{W}^\nabla, \overline{\nabla}).$$

In the case when $\mathcal{L} = \Theta_X, \mathcal{H} = \Theta_S$ the functor $F_!$ equals the functor introduced by Katz in [12].

Consider the LR-algebra $\mathcal{K} \rightarrow \Theta_X$ with the obvious anchor map. We may consider its generalized sheaf of enveloping algebras $\mathcal{U}(\mathcal{K})$ which is a sheaf of $\mathcal{O}_X$ -algebras, and a flat $\mathcal{L}$ -connection $(\mathcal{W}, \nabla)$ gives in a natural way a flat $\mathcal{K}$ -connection. We get in a natural way a left $\mathcal{U}(\mathcal{K})$ -module structure on $\mathcal{W}$. We may consider the category of sheaves of left $\mathcal{U}(\mathcal{K})$ -modules which has enough injectives. We can thus apply the machinery of derived functors. Let $\mathcal{U}, \mathcal{V}$ be two left $\mathcal{U}(\mathcal{K})$ -modules and consider the sheaf $\underline{\text{Hom}}_{\mathcal{U}(\mathcal{K})}(\mathcal{U}, \mathcal{V})$ which in a natural way is a left $\mathcal{U}(\mathcal{K})$ -module. There is by definition an equality

$$\underline{\text{Hom}}_{\mathcal{U}(\mathcal{K})}(\mathcal{U}, \mathcal{V})^\nabla = \underline{\text{Hom}}_{\mathcal{U}(\mathcal{K})}(\mathcal{U}, \mathcal{V})$$

hence by Proposition 1.6 it follows that $\underline{\text{Hom}}_{\mathcal{U}(\mathcal{K})}(\mathcal{U}, \mathcal{V})$ is a left $\mathcal{U}(\mathcal{H})$ -module, hence we get a left exact functor

$$\underline{\text{Hom}}_{\mathcal{U}(\mathcal{K})}(\mathcal{U}, -) : \underline{\mathcal{L}\text{-conn}} \rightarrow \underline{\mathcal{H}\text{-conn}}.$$

Definition 1.7. Let $\mathcal{U}, \mathcal{V}$ be left $\mathcal{U}(\mathcal{K})$ -modules. We define for all $i \geq 0$ the cohomology-sheaves

$$\underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^i(\mathcal{U}, \mathcal{V}) = R^i \underline{\text{Hom}}_{\mathcal{U}(\mathcal{K})}(\mathcal{U}, \mathcal{V}).$$

We write

$$\underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^*(\mathcal{U}, \mathcal{V}) = \bigoplus_{i \geq 0} \underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^i(\mathcal{U}, \mathcal{V}).$$

By construction there is for all $i \geq 0$ a canonical flat connection

$$\nabla_i : \mathcal{H} \rightarrow \underline{\text{End}}_{\mathcal{O}_T}(\underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^i(\mathcal{U}, \mathcal{V})).$$

We get thus a flat connection

$$\nabla : \mathcal{H} \rightarrow \underline{\text{End}}_{\mathcal{O}_T}(\underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^*(\mathcal{U}, \mathcal{V})),$$

called the *generalized Gauss-Manin connection*. We write

$$\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, \mathcal{V}) = \underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(\mathcal{O}_X, \mathcal{V}).$$

Proposition 1.8. *Assume $U = \mathrm{Spec}(A), V = \mathrm{Spec}(S)$ are open affine subsets of X and S with $f(U) \subseteq V$ and $\mathcal{K}$ a locally free LR-algebra. There is an isomorphism*

$$\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, \mathcal{W})|_U \cong \mathbf{H}^i(K, W)$$

where $\mathcal{K}|_U = K$ and $\mathcal{W}|_U = W$.

Proof. This is straightforward: By Proposition 1.5 we get isomorphisms

$$\begin{aligned} \underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, \mathcal{W})|_U &= \mathrm{R}^i \underline{\mathrm{Hom}}_{\mathcal{U}(\mathcal{K})}(\mathcal{O}_X, \mathcal{W})|_U = \mathrm{R}^i \mathrm{Hom}_{U(K)}(A, W) = \\ &= \mathrm{Ext}_{U(K)}^i(A, W) = \mathbf{H}^i(K, W) \end{aligned}$$

and the claim follows. $\square$

Hence the cohomology-sheaves $\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{L})}^i(X, \mathcal{W})$ may be calculated using the standard complex when $\mathcal{L}$ is locally free. We get in a natural way flat $\mathcal{H}$ -connections $\nabla_i = \mathrm{R}^i F(\nabla)$ on the cohomology sheaves $\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, \mathcal{W})$ - generalized Gauss-Manin connections - for all $i \geq 0$.

Notice that the quasi-coherent $\mathcal{O}_S$ -modules

$$\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, \mathcal{W})$$

specialize to $\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{L})}^i(X, \mathcal{W})$ when $S = T$. The cohomology sheaves $\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, \mathcal{W})$ generalize simultaneously several cohomology theories as mentioned in [17]. In particular if $\mathcal{L} = \Theta_X, \mathcal{H} = \Theta_S$ there are isomorphisms

$$\underline{\mathrm{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, \mathcal{W}) \cong \mathbf{H}^i(X/S, \mathcal{W}, \nabla)$$

where $\mathbf{H}^i(X/S, \mathcal{W}, \nabla)$ is the de Rham-cohomology sheaf introduced in 3.1 [12]. The generalized Gauss-Manin connection introduced in this section specialize to the Gauss-Manin connection introduced in [12] and [14].

Example 1.9. *Let $f : X = \mathrm{Spec}(B) \rightarrow S = \mathrm{Spec}(R)$ be a map of schemes and let W be a B -module with a flat connection*

$$\nabla : T_X \rightarrow \mathrm{End}_R(W).$$

Assume $L \subseteq T_X$ is a locally free LR-algebra, and consider the sub- R -algebra

$$B^L \subseteq B.$$

We get a map of schemes

$$\pi : X \rightarrow X/L = \mathrm{Spec}(B^L) \rightarrow S$$

and an associated exact sequence

$$0 \rightarrow \mathrm{Der}_{B^L}(B) \rightarrow \mathrm{Der}_R(B) \rightarrow B \otimes_{B^L} \mathrm{Der}_R(B^L)$$

Assume $H \subseteq \mathrm{Der}_R(B^L)$ is a sub-Lie algebra with $T_X \rightarrow \pi^* H$ surjective. We get thus a right exact sequence

$$0 \rightarrow L \rightarrow T_X \rightarrow \pi^* H \rightarrow 0$$

hence we get a functor

$$F_! : \underline{T_X - conn} \rightarrow \underline{H - conn}$$

as defined above. We get thus for all locally free Lie-algebras L canonical flat connections on the A^L -modules $\mathbf{H}^i(L, W)$

$$\nabla_i : H \rightarrow \text{End}_R(\mathbf{H}^i(L, W))$$

where (W, ∇) is a flat T_X -connection.

We end this section with an analogue of a proposition found in [12].

Proposition 1.10. *Suppose X is proper and smooth over S and k is a field of characteristic zero. Then for any flat T_X -connection $\mathcal{W}$ the sheaf $\underline{\text{Ext}}_{\mathcal{U}(T_f)}^i(X, \mathcal{W})$ is a locally free finite rank $\mathcal{O}_S$ -module.*

Proof. We have an isomorphism

$$\underline{\text{Ext}}_{\mathcal{U}(T_f)}^i(X, \mathcal{W}) \cong \mathbf{H}^i(X/S, \mathcal{W}, \nabla)$$

hence since by properness $\underline{\text{Ext}}_{\mathcal{U}(T_f)}^i(X, \mathcal{W})$ is a coherent $\mathcal{O}_S$ -module the statement follows from Proposition 8.8 [12]. $\square$

Note: many constructions on the Ext-functor introduced in [3] globalize to give similar results (Kunneth-formulas etc) for the $\underline{\text{Ext}}$ -functor introduced in this section. There is similarly defined a Tor-functor (up to non-canonical isomorphism) and constructions introduced in [3] globalize to give change of rings-spectral sequences.

2. APPLICATION: A GYSIN MAP FOR K-THEORY OF FLAT CONNECTIONS

In this section we prove existence of a Gysin-map map between K-groups of categories of flat connections.

Let in the following $\underline{\mathcal{C}}, \underline{\mathcal{D}}$ be two arbitrary abelian categories with $\underline{\mathcal{C}}$ having enough injectives and let $F : \underline{\mathcal{C}} \rightarrow \underline{\mathcal{D}}$ be a left exact functor with the property that $R^i F(C) = 0$ for all objects C with $i \gg 0$. We may thus construct the right derived functors $R^i F$ of F (see [7]) having the property that for any exact sequence

$$0 \rightarrow C' \rightarrow C \rightarrow C'' \rightarrow 0$$

in $\underline{\mathcal{C}}$, we get a long exact sequence of objects in $\underline{\mathcal{D}}$:

$$\begin{aligned} 0 \rightarrow F(C') \rightarrow F(C) \rightarrow F(C'') \rightarrow \\ R^1 F(C') \rightarrow R^1 F(C) \rightarrow R^1 F(C'') \rightarrow \dots \end{aligned}$$

It follows we get an induced push-forward map at the level of K-groups as follows: Let $K(\underline{\mathcal{C}})$ and $K(\underline{\mathcal{D}})$ be the K-group of the categories $\underline{\mathcal{C}}$ and $\underline{\mathcal{D}}$, ie $K(\underline{\mathcal{C}})$ is the free abelian group on the isomorphism classes in $\underline{\mathcal{C}}$ modulo the subgroup generated by elements of the type

$$[C] - [C'] - [C'']$$

arising from an exact sequence in $\underline{\mathcal{C}}$ as usual.

Definition 2.1. Let C be an object of $\underline{\mathcal{C}}$. Make the following definition:

$$F_! [C] = \sum_{i \geq 0} (-1)^i [R^i F(C)].$$

Proposition 2.2. *Let $F : \underline{\mathcal{C}} \rightarrow \underline{\mathcal{D}}$ be a left exact functor. The map $F_!$ induces a well-defined map*

$$F_! : K(\underline{\mathcal{C}}) \rightarrow K(\underline{\mathcal{D}})$$

at the level of K-groups.

Proof. This is straightforward. $\square$

Let $K(X)$ (resp. $K(S)$) be the K-group of the category of coherent sheaves on a scheme X (resp. S) and let furthermore $f : X \rightarrow S$ be a proper map of schemes.

Corollary 2.3. *If $F = f_*$ is the pushforward map arising from a map of schemes $f : X \rightarrow S$ the induced map $F_! = f_!$ at level of K-groups is the Gysin map*

$$f_! : K(X) \rightarrow K(S),$$

defined by

$$f_![\mathcal{E}] = \sum_{i \geq 0} (-1)^i [R^i f_* \mathcal{E}].$$

Proof. See [7]. $\square$

Let $(F, f) : (\mathcal{L}, X) \rightarrow (\mathcal{H}, S)$ be a surjective morphism of LR-algebras, where $f : X \rightarrow S$ is a map of schemes over a scheme T . Let furthermore

$$\underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, -) : \underline{\mathcal{L} - \text{conn}} \rightarrow \underline{\mathcal{H} - \text{conn}}$$

be the Ext-sheaves from Definition 1.7.

We get thus for any exact sequence of flat connections in $\underline{\mathcal{L} - \text{conn}}$ a long exact sequence in $\underline{\mathcal{H} - \text{conn}}$, hence we get an induced map at level of K-theory: Let $K(\mathcal{L})$ (resp. $K(\mathcal{H})$) be the K-groups of the categories of coherent sheaves of flat $\mathcal{L}$ -connections (resp. $\mathcal{H}$ -connections). We can prove existence of a *Gysin-map* between these K-groups. Assume for simplicity for the rest of this section that $\underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, W) = 0$ for all W and all $i \gg 0$ (this holds for instance in the situation considered in Proposition 1.10).

Theorem 2.4. *There is a Gysin map*

$$F_! : K(\mathcal{L}) \rightarrow K(\mathcal{H})$$

defined as follows:

$$F_![\mathcal{W}, \nabla] = \sum_{i \geq 0} (-1)^i [\underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^i(X, \mathcal{W})].$$

Proof. The claim of the theorem follows from general properties of the Ext-functor, Proposition 2.2 as follows: For an exact sequence

$$0 \rightarrow (\mathcal{W}', \nabla') \rightarrow (\mathcal{W}, \nabla) \rightarrow (\mathcal{W}'', \nabla'') \rightarrow 0$$

of horizontal maps of flat $\mathcal{L}$ -connections, we always get by the functorial properties of Ext a long exact sequence

$$\begin{aligned} 0 \rightarrow \underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^0(X, \mathcal{W}') \rightarrow \underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^0(X, \mathcal{W}) \rightarrow \underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^0(X, \mathcal{W}'') \rightarrow \\ \underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^1(X, \mathcal{W}') \rightarrow \underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^1(X, \mathcal{W}) \rightarrow \underline{\text{Ext}}_{\mathcal{U}(\mathcal{K})}^1(X, \mathcal{W}'') \rightarrow \dots \end{aligned}$$

of $\mathcal{H}$ -connections hence the map is well-defined. $\square$

Assume $X = \text{Spec}(B)$, $S = \text{Spec}(A)$ and $\mathcal{K} = K$ where K is locally free. We get in this case a Gysin map

$$F_! : K(L) \rightarrow K(H)$$

defined by

$$F_!(W, \nabla) = \sum_{i \geq 0} (-1)^i [\mathbf{H}^i(K, W)]$$

hence the standard complex calculates the Gysin-map in the case when K is locally free.

In the special case when $\mathcal{L} = \Theta_X$ and $\mathcal{H} = \Theta_S$ we get a corollary:

Corollary 2.5. *Assume the base field is of characteristic zero and that X is proper over S . There is a Gysin-map*

$$F_! : K(\Theta_X) \rightarrow K(\Theta_S)$$

defined by

$$F_!(\mathcal{W}, \nabla) = \sum_{i \geq 0} (-1)^i \mathbf{H}^i(X/S, \mathcal{W}, \nabla),$$

where $\mathbf{H}^i(X/S, \mathcal{W}, \nabla)$ is the group introduced in 3.1,[12].

Proof. This follows from Proposition 1.10. $\square$

There is ongoing work proving a Riemann-Roch formula using the Gysin-map introduced in this note and a refined version of the Chern-character introduced in [18].

3. EXPLICIT EXAMPLES: A GAUSS-MANIN CONNECTION FOR A FAMILY OF CURVES

We consider examples of Gauss-Manin connections on a Lie-Rinehart algebra $H \subseteq T_C$ on a family of curves $X \rightarrow C$, using connections constructed in previous papers on the subject (see [16] and [17] for details on the calculations).

Consider the polynomial $B_{mn2} = x^m + y^n + z^2$ and let $A = \mathbf{C}[x, y, z]/x^m + y^n + z^2$ where $\mathbf{C}$ is the complex numbers. By the results of [17] we get a family of maximal Cohen-Macaulay modules on A given by the two matrices

$$\phi = \begin{pmatrix} x^{m-k} & y^{n-l} & 0 & z \\ y^l & -x^k & z & 0 \\ z & 0 & -y^{n-l} & -x^k \\ 0 & z & x^{m-k} & -y^l \end{pmatrix}$$

and

$$\psi = \begin{pmatrix} x^k & y^{n-l} & z & 0 \\ y^l & -x^{m-k} & 0 & z \\ 0 & z & -y^l & x^k \\ z & 0 & -x^{m-k} & -y^{n-l} \end{pmatrix},$$

where $1 \leq k \leq m$ and $1 \leq l \leq n$. The matrices ϕ and ψ gives rise to an exact sequence of A -modules

$$(3.0.1) \quad \dots \rightarrow^\psi P \rightarrow^\phi P \rightarrow^\psi P \rightarrow^\phi P \rightarrow W(\phi, \psi) \rightarrow 0.$$

And by the results in [6] the module $W = W(\phi, \psi)$ is a maximal Cohen-Macaulay module on A . The pair (ϕ, ψ) is a *matrix-factorization* of the polynomial f . In [17] we compute explicitly algebraic $\mathbf{V}_W$ -connections on the modules $W = W(\phi, \psi)$ for all $1 \leq k \leq m$ and $1 \leq l \leq n$. The calculation is as follows: We compute generators and syzygies of the derivation-modules $\text{Der}_{\mathbf{C}}(A)$ and find it is generated by the derivations

$$\begin{aligned} \delta_0 &= 2nx\partial_x + 2my\partial_y + mnz\partial_z \\ \delta_1 &= mx^{m-1}\partial_y - ny^{n-1}\partial_z \\ \delta_2 &= -2z\partial_x + mx^{m-1}\partial_z \end{aligned}$$

$$\delta_3 = -2z\partial_y + ny^{n-1}\partial_z.$$

The syzygy-matrix of $\text{Der}_{\mathbf{C}}(A)$ is the following matrix

$$\rho = \begin{pmatrix} y^{n-1} & z & 0 & x^{m-1} \\ 2x & 0 & -2z & -2y \\ 0 & nx & ny^{n-1} & -nz \\ -mz & my & -mx^{m-1} & 0 \end{pmatrix}.$$

A calculation using the Kodaira-Spencer map $g : \text{Der}_{\mathbf{C}}(A) \rightarrow \text{Ext}_A^1(W, W)$ gives the following connection:

$$(3.0.2) \quad \overline{\nabla}_{\delta_0} = \delta_0 + A_0 = \delta_0 + \begin{pmatrix} nk + ml - \frac{1}{2}mn & 0 & 0 & 0 \\ 0 & \frac{3}{2}mn - ml - nk & 0 & 0 \\ 0 & 0 & \frac{1}{2}mn + ml - nk & 0 \\ 0 & 0 & 0 & \frac{1}{2}mn + nk - ml \end{pmatrix}.$$

$$(3.0.3) \quad \overline{\nabla}_{\delta_1} = \delta_1 + \begin{pmatrix} 0 & b_2 & 0 & 0 \\ b_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & b_4 \\ 0 & 0 & b_3 & 0 \end{pmatrix} = \delta_1 + A_1,$$

with $b_1 = \frac{1}{4}(mn - 2nk - 2ml)x^{k-1}y^{l-1}$, $b_2 = \frac{1}{4}(3mn - 2ml - 2nk)x^{m-k-1}y^{n-l-1}$, $b_3 = \frac{1}{4}(2nk - mn - 2ml)x^{m-k-1}y^{l-1}$ and $b_4 = \frac{1}{4}(2nk - 2ml + mn)x^{k-1}y^{n-l-1}$.

$$(3.0.4) \quad \overline{\nabla}_{\delta_2} = \delta_2 + \begin{pmatrix} 0 & 0 & c_3 & 0 \\ 0 & 0 & 0 & c_4 \\ c_1 & 0 & 0 & 0 \\ 0 & c_2 & 0 & 0 \end{pmatrix} = \delta_2 + A_2,$$

with $c_1 = \frac{1}{n}(\frac{1}{2}mn - ml - nk)x^{k-1}$, $c_2 = \frac{1}{n}(\frac{3}{2}mn - ml - nk)x^{m-k-1}$, $c_3 = \frac{1}{n}(\frac{1}{2}mn + ml - nk)x^{m-k-1}$ and $c_4 = \frac{1}{n}(ml - nk - \frac{1}{2}mn)x^{k-1}$.

$$(3.0.5) \quad \overline{\nabla}_{\delta_3} = \delta_3 + \begin{pmatrix} 0 & 0 & 0 & d_4 \\ 0 & 0 & d_3 & 0 \\ 0 & d_2 & 0 & 0 \\ d_1 & 0 & 0 & 0 \end{pmatrix} = \delta_3 + A_3,$$

where $d_1 = \frac{1}{m}(\frac{1}{2}mn - ml - nk)y^{l-1}$, $d_2 = \frac{1}{m}(ml + nk - \frac{3}{2}mn)y^{n-l-1}$, $d_3 = \frac{1}{m}(\frac{1}{2}mn + ml - nk)y^{l-1}$ and $d_4 = \frac{1}{m}(\frac{1}{2}mn - ml + nk)y^{n-l-1}$.

Theorem 3.1. *For all $1 \leq k \leq m$ and $1 \leq l \leq n$ the equations 3.0.2-3.0.5 define algebraic $\mathbf{V}_W$ -connections $\overline{\nabla}^{\phi, \psi} : \mathbf{V}_W \rightarrow \text{End}_{\mathbf{C}}(W)$ where $W = W(\phi, \psi)$.*

Proof. See Theorem 2.1 in [17] □

Note that the vectorfield δ_0 - the Euler derivation - generates a locally free Lie-Rinehart algebra $L \subseteq T_X$. Let $X = \text{Spec}(A)$ and $C = \text{Spec}(A^L)$ with associated map of schemes

$$\pi : X \rightarrow C.$$

From Example 1.9 we get a right exact sequence

$$L \rightarrow T_X \rightarrow \pi^*H \rightarrow 0$$

where $H \subseteq T_C$ is a sub-Lie algebra, hence we get a functor

$$\underline{T_X - conn} \rightarrow \underline{H - conn}.$$

We get thus connections

$$\nabla_i : H \rightarrow \text{End}(\mathbf{H}^i(L, W))$$

for all $i \geq 0$.

Theorem 3.2. *There exists for all $1 \leq k \leq m$ and $1 \leq l \leq n$ a Gauss-Manin connection*

$$\nabla_i : H \rightarrow \text{End}(\mathbf{H}^i(L, W))$$

for all $i \geq 0$.

Proof. This is immediate from the discussion above and the results in section one of this paper. $\square$

We illustrate the theory with some basic examples.

Example 3.3. *Consider the following situation: Let $F = \mathbf{C}$ be the field of complex numbers, $A = F[x, y]$ and $X = \text{Spec}(A)$. Let L be the Lie-algebra generated by $\partial_x + \partial_y \in \text{Der}_F(A)$ and $S = \text{Spec}(A^L) = \text{Spec}(F[x - y]) = \text{Spec}(F[T])$. We have a map of schemes $\pi : X \rightarrow S$ and there is an exact sequence*

$$0 \rightarrow L \rightarrow T_X \rightarrow \pi^* T_S \rightarrow 0$$

of sheaves. Let E be the free A -module of rank two. Then E has a flat T_X -connection

$$\nabla : T_X \rightarrow \text{End}_F(E).$$

A calculation shows that $\mathbf{H}^0(L, E)$ is the $F[T]$ -module

$$E^L = A^L \oplus A^L = F[T] \oplus F[T]$$

and $\mathbf{H}^1(L, E) = 0$. The Gauss-Manin-connection on

$$\mathbf{H}^0(L, E) = E^L = F[T] \oplus F[T]$$

is the connection

$$\nabla : T_S \rightarrow \text{End}_F(\mathbf{H}^0(L, E))$$

given by

$$\nabla(\partial)(a, b) = (\partial(a), \partial(b))$$

where $\partial = \frac{\partial}{\partial T}$.

Example 3.4. *Consider the map*

$$\pi : S \rightarrow C$$

where $S = \text{Spec}(F[x, y])$ and $C = \text{Spec}(F[t])$ defined by

$$\pi(t) = x^m - y^n = f(x, y).$$

We get a relative tangent sequence

$$0 \rightarrow T(\pi) \rightarrow T_S \rightarrow \pi^* T_C$$

which is not right exact, since the map π is not smooth. We calculate, and find $T(\pi)$ to be defined by the vector field

$$\partial = mx^{m-1}\partial_x + ny^{n-1}\partial_y.$$

The vector field ∂ and the Euler-vectorfield

$$E = nx\partial_x + my\partial_y$$

defines a Lie-sub-algebra

$$L = \{\partial, E\} \subseteq \text{Der}_F(B)$$

where $B = F[x, y]$, and we check we get an exact sequence

$$0 \rightarrow T(\pi) \rightarrow L \rightarrow \pi^*H \rightarrow 0$$

where $H = \text{mnt}\partial_t \subseteq F[t]\partial_t = T_C$. We get thus for any flat L -connection W a flat H -connection

$$\nabla : H \rightarrow \text{End}_F(\mathbf{H}^i(T(\pi), W)),$$

a logarithmic Gauss-Manin connection. The singular fibre $\pi^{-1}(0) = \text{Spec}(F[x, y]/x^m - y^n)$ of the map π is defined by the ideal $I = (f) = (x^m - y^n)$, and there is an obvious flat $T(\pi)$ -connection

$$\nabla : T(\pi) \rightarrow \text{End}_F(I)$$

defined by

$$\nabla(x)(b) = x(b).$$

We thus get natural flat H -connections

$$\nabla_i : H \rightarrow \text{End}_F(\mathbf{H}^i(T(\pi), I))$$

for $i = 0, 1$, and we aim to calculate ∇_i . A calculation shows that

$$\mathbf{H}^0(T(\pi), I) = I^{T(\pi)} = F[t].$$

The standard complex $C^p(T(\pi), I)$ looks as follows:

$$I \rightarrow \text{Hom}_B(T(\pi), I)$$

where $B = F[x, y]$. It follows that $T(\pi) = B\partial$ and $I = (f)$. Hence

$$\text{Hom}_B(T(\pi), I) = Bf.$$

We get thus the complex

$$\partial : Bf \rightarrow Bf$$

and

$$\mathbf{H}^1(T(\pi), I) = Bf/\text{Im}(\partial).$$

A calculation shows that if $m \leq n$

$$\dim(H^1(T(\pi), I)) = \frac{(n-2)(m-1)}{2}.$$

There is ongoing work implementing computer algorithms for calculations of Gauss-Manin connections.

Acknowledgements: Thanks to Paul Bressler, Sophie Chemla, Christopher Deninger, Nicholas Katz and Max Karoubi for email correspondence, discussions and comments on the contents of this paper.

REFERENCES

- [1] R. Almeida, A. Kumpera, Structure produit dans la categorie des algebroides de Lie, *Ann. Acad. Brasil. Cienc.* 53, 247-250 (1981)
- [2] J. Bernstein, D-modules and functors, *manuscript*
- [3] H. Cartan, S. Eilenberg, Homological algebra, *Princeton Math. Series* (1956)
- [4] S. Bloch, H. Esnault, A Riemann-Roch theorem for flat bundles with values in algebraic Chern-Simons theory, *Annals of Mathematics* 151 (2000)
- [5] S. Chemla, A duality property for complex Lie-algebroids, *Math. Zeitschrift*, 232 (1999)
- [6] D. Eisenbud, Homological algebra on a complete intersection with an application to group representations, *Trans. Amer. Math. Soc.* vol. 260 (1980)
- [7] R. Hartshorne, Algebraic geometry, Graduate Texts in Mathematics no. 52, *Springer Verlag* (1977)
- [8] J. Huebschmann, Poisson cohomology and quantization, *J. für die reine und angewandte Mathematik* vol. 408 (1990)
- [9] F. W. Kamber, P. Tondeur, Characteristic invariants of foliated bundles, *Manuscr. Math.* 11 (1974)
- [10] F. W. Kamber, P. Tondeur, Invariant differential operators and the cohomology of Lie-algebra sheaves, *Mem. of the AMS* no. 113 (1971)
- [11] M. Karoubi, Classes caracteristiques de fibres feuilletes, holomorphes ou algebriques, *K-theory* 8 (1994)
- [12] N. Katz, Nilpotent connections and the monodromy theorem, *Publ. Math. IHES* 39 (1970)
- [13] N. Katz, On the differential equation satisfied by period matrices, *Publ. Math. IHES* 35 (1968)
- [14] N. Katz, T. Oda, On the differentiation of De Rham cohomology classes with respect to parameters, *J. Math. Kyoto Univ.* 8 (1968)
- [15] H. Maakestad, A universal Riemann-Roch formula, *In progress*
- [16] H. Maakestad, Chern-klasser for maximale Cohen-Macaulay moduler paa normale kvotient singulariteter, *Master Thesis*, University of Oslo (1995)
- [17] H. Maakestad, Chern-classes and Lie-Rinehart algebras, *math.AG/0306254*
- [18] H. Maakestad, The Chern-character for Lie-Rinehart algebras, *Ann. Inst. Fourier* vol. 55 (2005) Fasc. 7
- [19] N. Saavedra Rivano, Categories Tannakiennes, *Lecture Notes in Mathematics* Vol. 265

INSTITUT DE MATHÉMATIQUES, UNIVERSITÉ PARIS VII
 E-mail address: `maakestad@math.jussieu.fr`